

Published on *SeniorNavigator* (<https://seniornavigator.org>)

[5 Things to Know About Taking 5 or More Medications](#)

Taking multiple drugs increases your risk for falls, confusion and other serious side effects

[EN ESPAÑOL](#)

Medicines can improve our lives — and save them — by regulating blood pressure, curing infections and calming restless minds and aching joints. But sometimes, too much of a good thing isn't very good at all.

More than 40 percent of older Americans regularly take five or more prescription drugs, and nearly 20 percent take 10 or more, according to [a 2020 report](#) from the nonpartisan think tank Lown Institute. When over-the-counter medicines and supplements are factored in, the share of older adults popping five or more pills — a practice known as polypharmacy — shoots up to 67 percent.

"People can get into this situation quite easily," says Lon Schneider, M.D., a geriatric psychiatrist and professor of psychiatry, neurology and gerontology at the University of Southern California, Los Angeles.

For example, a primary care physician may treat a patient for [high blood pressure](#) with medication that can result in more frequent urination. At the same time, that patient may visit a urologist for incontinence "and get a medication that can further affect blood pressure and cognition, as well," Schneider says. "They go to a gastroenterologist because of indigestion and get something for that, and then the primary care [doctor] — without knowing any of this — may give them something for anxiety and depression and something else for sleep. It adds up very quickly."

And this medication overload can have serious consequences: Every day, 750 older adults are hospitalized for side effects from medication, the Lown Institute report states. If current trends continue, the problem could result in more than 4.5 million hospitalizations and 150,000 premature deaths among older adults over the next decade.

With increased age comes increased risk

The incidence of [chronic disease increases with age](#). Nearly 80 percent of adults 55 and older have at least one chronic condition, data from the Centers for Disease and Prevention (CDC) shows; almost half have two or more. And hand in hand with these conditions come medications to manage them.

The problem is that as we age, our bodies are less able to handle some drugs. "Medications that are completely fine for people in their 20s and 40s may not be fine for people over 60," says Jeffrey Keller, director of the Institute for Dementia Research and Prevention in Baton Rouge, Louisiana.

One reason: Aging alters the body's capacity to absorb, metabolize and excrete drugs, Schneider says. "So you're often getting relatively more of that drug than you may need, especially when you reach 65 to 70. The dose you took earlier in your life with no problem is now actually an increased dose."

What's more, clinical trials that test the safety of approved drugs "are conducted in a way that minimizes potential interactions with other medications," Keller says. "However, in real life, there are very real interactions for older people that were not observed in the trials."

Drug interactions can cause dangerous side effects

The number of potentially dangerous drug-drug interactions is impossible to estimate, because the potential polypharmacy combinations are practically limitless. One example of a combination to avoid is blood thinners, such as warfarin, and nonsteroidal anti-inflammatory drugs (NSAIDs), including ibuprofen (Advil) or naproxen (Aleve). Taking these two together can increase bleeding risks.

5 most commonly used prescription drugs

Use among U.S. adults ages 60 to 79

- Lipid-lowering drugs (45 percent)
- Antidiabetic agents (23.6 percent)
- Beta blockers (for high blood pressure or heart disease, 22.3 percent)
- ACE inhibitors (21.3 percent)
- Proton pump inhibitors (16.9 percent)

Source: CDC, 2019/National Health and Nutrition Examination Survey

For older adults, cognitive problems like memory issues, disorientation, confusion and clouded thinking are a top concern of taking multiple medications, Schneider says. So are [falls](#). And research shows that the more medications a person takes, the greater the risk for falling.

"A hip fracture in someone in their 70s or 80s is really quite serious and has major implications and effects on mortality," Schneider says. In fact, falls double the risk of death for an elderly person in the first year after the break, one study shows; the risk remains close to double for several years later.

Polypharmacy can affect brain health

While your daily drug regimen may be designed to improve your joint health or mental health, it could negatively affect your brain health.

For starters, several studies have linked polypharmacy to dementia. What's more, commonly prescribed drugs — from antidepressants to corticosteroids — can cause cognitive problems that mimic dementia. However, unlike dementia, these symptoms are often curable, which is why Keller always considers medications when a person is concerned about their brain health.

"When someone comes in with suspected cognitive decline, the first thing we do is try to find a reversible cause," he says.

Some classes of medications also have been associated with increased risk for dementia, like benzodiazepines, prescribed to treat conditions from anxiety to insomnia, and [anticholinergics](#), which are used to treat a range of conditions including chronic obstructive pulmonary disease and gastrointestinal disorders. The over-the-counter allergy medicine diphenhydramine (Benadryl) also has anticholinergic effects.

A 2019 study published in JAMA Internal Medicine found that longer-term use of anticholinergics was associated with about a 50 percent higher dementia risk, compared to short-term use.

It's not completely clear how the drugs might increase dementia risk, but a 2016 study found that the brains of those on long-term anticholinergics had atrophied and the hollow spaces inside increased. The neurons also had reduced ability to metabolize glucose. The higher the drug exposure, the more pronounced the effect.

Anticholinergics can also counteract the medications given to improve memory and thought in Alzheimer's disease, making a bad situation even worse. Three common Alzheimer's drugs (brand names Aricept, Exelon and Razadyne) belong to a class called cholinesterase inhibitors. They work by increasing acetylcholine, a chemical that allows the body's nerve cells to communicate with each other, by preventing its breakdown.

Anticholinergics do the opposite. They block the ability of acetylcholine to exert its effect, lowering nerve signaling, thereby decreasing whatever activity those nerves control.

Other hidden dangers lurk in common medicines

Cognitive changes aren't the only potential side effects of these medications, said Shanna Trenaman, a pharmacist at Dalhousie University in Halifax, Nova Scotia, who specializes in geriatrics. Plenty of studies have found that anticholinergics can increase stomach upset and cause constipation and urinary retention.

Some sleep aids (including a number that are sold over the counter) also have anticholinergic effects; these have been linked to increased risk of car wrecks, falls

and other injuries, Trenaman says. And medication interactions are possible with these, as well.

"I would encourage everyone to think critically about whether or not they need to take medication for sleep," Trenaman told AARP. "They help you fall asleep only about seven minutes faster on average, so is that worth these risks?"

If you are taking something to help you sleep and decide to discontinue it, do it with guidance from a health care professional who can help you taper off the medication slowly.

A health care provider should review your medications often

Experts interviewed for this article have a unanimous recommendation: Be proactive in your medication decisions.

Older adults managing multiple health conditions may see several doctors — and each may be unaware of what the others have prescribed. Recently there's been greater attention focused on the concept of deprescribing — or "evaluating people's medications, consciously assessing whether it's really needed, and planning how best to reduce and discontinue it," Schneider says. It's important to note, however, that patients should never stop taking a drug without consulting a doctor, as this can be dangerous.

Instead, the three experts recommend at least an annual review of medications with your main physician. Bring all of your medications in — don't forget about any over-the-counter medicines you are taking, plus vitamins and supplements — and talk about potential interactions and which drugs you may no longer need.

"There's a perception that all over-the-counter medications are as safe as water, but that's just not true. Antihistamines, anticholinergics and sleep aids are just as potent over the counter as they are by prescription, and they are big contributors to this issue," Schneider says.

Your pharmacist can also help answer any medication questions, especially when it comes to spotting potentially troublesome drug combinations.

AARP was founded in 1958 and has over 38 million members. It is a nonprofit, nonpartisan organization for people over the age of 50. AARP is well-known for its advocacy efforts, providing its members with important information, products and services that enhance quality of life as they age. They also promote community service and keep members and the public informed on issues relating to the over 50 age group.

Article Source

AARP

Source URL

<https://www.aarp.org>

Last Reviewed

Thursday, April 4, 2024